

L7 ANSWER 4 OF 5 MEDLINE MEDLINE DUPLICATE 3
 ACCESSION NUMBER: 2000013158 MEDLINE
 DOCUMENT NUMBER: 20013158 PubMed ID: 10544301
 TITLE: Overexpression of the **EphA2** tyrosine kinase in prostate cancer.
 AUTHOR: Walker-Daniels J; Coffman K; Azimi M; Rhim J S; Bostwick D G; Snyder P; Kerns B J; Waters D J; Kinch M S
 CORPORATE SOURCE: Department of Basic Medical Sciences, Purdue University, West Lafayette, Indiana 47907-1246, USA.
 SOURCE: PROSTATE, (1999 Dec 1) 41 (4) 275-80.
 Journal code: 8101368. ISSN: 0270-4137.
 PUB. COUNTRY: United States
 DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)
 LANGUAGE: English
 FILE SEGMENT: Priority Journals
 ENTRY MONTH: 199912
 ENTRY DATE: Entered STN: 20000113
 Last Updated on STN: 20000113
 Entered Medline: 19991222

AB BACKGROUND: Molecules that are highly expressed by human prostate cancers may serve as therapeutically relevant targets or tumor markers. Tyrosine kinases are frequently overexpressed in **metastatic** tumor cells and this prompted us to screen for tyrosine kinases that are overexpressed in prostate cancer cells. METHODS: Expression levels of the **EphA2** receptor tyrosine kinase were determined by Western blot analysis in canine and human prostate cancer cell lines and in immortalized and transformed variants of 267B1 prostatic epithelial cells. **EphA2** levels in benign human prostate and prostate cancers were also determined in formalin-fixed, paraffin-embedded tissues using immunohistochemical staining. RESULTS: **Metastatic** prostate cancer cells overexpressed **EphA2** by 10-100 fold as compared with non-invasive prostatic epithelial cells. **EphA2** immunoreactivity in vivo was also significantly greater in human prostate cancers as compared with benign prostate epithelium. CONCLUSIONS: The **EphA2** receptor tyrosine kinase is differentially expressed in human and canine prostate cancer cell lines and overexpressed in human prostate cancers as compared with benign prostate tissues. **Metastasis**-derived canine prostate carcinoma cell lines overexpress **EphA2** and may provide pre-clinical models to further evaluate the role of **EphA2** in prostate carcinogenesis. Further investigations are needed to determine the utility of **EphA2** as a tumor marker and a novel target in human prostate cancer.
 Copyright 1999 Wiley-Liss, Inc.